


William A. Tyndall, M.D., Ph.D. 3000 Fairway Drive, Altoona, PA 16602 814-231-2101

Routine For: Created By: DR TYNDALL May 26, 2003

HIP / KNEE - 64 Antiemboli Isometric




Extending toes toward knee, tense muscles on front of thigh and simultaneously squeeze buttocks. Keep leg and buttock flat on floor. Hold <u>6</u> seconds.

Repeat 20 times per set. Do 2 sets per session.

Do 3 sessions per day.


HIP / KNEE - 17 Strengthening: Straight Leg Raise, Phase I



Tighten muscle on front of thigh, then lift leg 8-10 inches from surface, keeping knee locked. Hold 6 seconds.

Repeat 20 times. Do 2 sessions per day.


HIP / KNEE - 11 Strengthening: Quadriceps Sets



Tighten muscles on top of thigh by pushing knees down into surface. Hold 6 seconds.

Repeat 20 times. Do 2 sessions per day.

HIP / KNEE - 18 Strengthening: Straight Leg Raise, Phase II



Resting on forearms, tighten muscle on front of thigh, then lift leg 8-10 inches from surface, keeping knee locked. Hold 6 seconds.

Repeat <u>20</u> times. Do <u>2</u> sessions per day.